Listing of the Claims

This listing of claims will replace all prior versions, and listings of claims in the application.

- (Previously presented) A method for treating a mammalian subject having a
 carcinoid tumor and carcinoid syndrome, comprising administering to the subject an amount of a
 therapeutic virus effective to treat the tumor and decrease one or more symptoms of the carcinoid
 syndrome, wherein the virus is a Newcastle Disease virus.
 - 2-4. (Cancelled).
- (Previously presented) The method of claim 1, wherein the virus is a mesogenic strain of Newcastle Disease Virus.
 - 6. (Original) The method of claim 1, wherein the virus is administered systemically.
 - 7. (Original) The method of claim 6, wherein the virus is administered intravenously.
- (Original) The method of claim 7, wherein the virus administered is a mesogenic strain of Newcastle Disease Virus.
 - 9-10. (Cancelled)
- 11. (Currently amended) The method of <u>elaim 1</u>, wherein the <u>virus is administered over an</u> administration time period-is of at least 1 hour.
- (Original) The method of claim 11, wherein the administration time period is at least 3 hours.
- 13. (Original) The method of claim 8, wherein the therapeutic virus is administered to the subject in one or more cycles, wherein at least one cycle comprises administering sequentially one or more desensitization doses of the virus followed by administering one or

more escalated doses of the virus, wherein the amount of the virus in each escalated dose is higher than the amount of virus in each desensitization dose.

- 14. (Original) The method of claim 13, wherein the cycle comprises one desensitization dose of from 1.2×10^{10} PFU to 4.8×10^{10} PFU per square meter of patient surface area, and one or more escalated doses of from 2.4×10^{10} PFU to 1.2×10^{11} PFU per square meter of patient surface area.
 - 15. (Cancelled)
 - 16. (Original) The method of claim 1, wherein the subject is a human subject.
 - 17. (Original) The method of claim 1, wherein the subject is a non-human mammal.
- 18. (Original) The method of claim 1, wherein the size of the tumor decreases after administration of the virus.
- (Original) The method of claim 1, wherein the level of 5-hydroxyindole acetic acid in urine of the subject decreases after administration of the virus.
 - 20. (Cancelled)
- 21. (Previously presented) The method of claim 1, wherein the one or more symptoms of carcinoid syndrome are selected from the group consisting of diarrhea, flushing, and fatigue.
 - 22-23. (Cancelled).